

**ACTIVATION OF SODIUM-HYDROGEN EXCHANGE IN HEART CELLS BY HYPEROSMOLAR SOLUTIONS.**

David Mhalley, MD, Paul Hamsworth, BSc, Halge Rasmussen, MD, Royal North Shore Hospital, Sydney, Australia.

Exposure of cardiac tissue to a hyperosmolar milieu is known to increase intracellular  $\text{Ca}^{2+}$  and resting tension. This enhances cellular acid production and may therefore be expected to decrease intracellular pH ( $\text{pH}_i$ ). Evidence from non-cardiac cells, however, indicates that hyperosmolality increases rather than decreases  $\text{pH}_i$ . We examined the effect of osmolality on  $\text{pH}_i$  in cardiac tissue. The resting membrane potential ( $E_m$ ) of guinea pig RV papillary muscles was recorded with conventional microelectrodes;  $\text{pH}_i$  was measured with ion-sensitive microelectrodes.  $E_m$  of 6 specimens in isotonic, modified HEPES-Tyrodé's solution (315 mOsm,  $\text{pH}$  7.40 at  $35^\circ\text{C}$ ) was  $83.7 \pm 1.7$  mV (mean  $\pm$  SE) while  $\text{pH}_i$  was  $7.09 \pm 0.02$ .  $E_m$  hyperpolarized by  $2.7 \pm 0.65$  mV ( $p < 0.01$ ) and  $\text{pH}_i$  increased by  $0.10 \pm 0.03$  ( $p < 0.02$ ) after switching to a similar superfusate rendered hyperosmolar (420 mOsm) with sucrose. In search of a mechanism for the alkaline shift we superfused 5 additional specimens with Tyrodé's solution containing  $5 \times 10^{-5}$  M 5-(N,N-dimethyl)amiloride (DMA). The  $\text{pH}_i$  in DMA containing solution was  $6.98 \pm 0.03$ .  $E_m$  hyperpolarized by  $3.1 \pm 1.1$  mV ( $p < 0.05$ ) and  $\text{pH}_i$  decreased by  $0.06 \pm 0.01$  units ( $p < 0.02$ ) after switching to DMA-containing hyperosmolar solution. A similar response to hyperosmolality was seen when extracellular  $\text{Na}^+$  was reduced to 15mM. We conclude that exposure to a hyperosmolar milieu causes an alkalization within heart cells. Abolition of this response by DMA and by a reduction of extracellular  $\text{Na}^+$  strongly suggests that it is mediated by activation of the sarcolemmal  $\text{Na}^+/\text{H}^+$  antiport.

**JUNCTIONAL ZONE BETWEEN SEGMENTS OF VARYING ACTION POTENTIAL DURATION APPEARS TO BE A SITE OF ARRHYTHMOGENESIS**

Zhen-Yuan Li, Ph.D., Takeshi Noda, M.D., Claudio Maldonado, M.S., Joel Kupersmith, M.D., F.A.C.C., University of Louisville, Louisville, KY.

We examined activity at the junctional zone between sheep Purkinje fiber ( $n=15$ ) displaying repolarization abnormalities and normal tissue in a double chamber bath using 3 standard microelectrodes. Ethylenediamine tetraacetate (3.3 mM) was added to one chamber (ABN). One microelectrode was located in ABN 3-4 mm from the junction. The other chamber, bathed in unmodified Tyrodé's, had a microelectrode at the junction (NL-J) and another 3-4 mm distally in normal tissue (NL-D). At 5-10 minutes: the action potential in ABN but not in NL was prolonged. A second plateau (at  $-65 \pm 3$  mV) ( $n=9$ ) or a gradually depolarizing slope resembling normal automaticity ( $n=6$ ) appeared at NL-J but not at NL-D. In both instances, two phenomena occurred: 1) There were spontaneous activations originating at NL-J and conducting to NL-D as normal action potentials, and to ABN as early afterdepolarizations; 2) Triggered activity in NL-D was easily induced by early afterdepolarizations from ABN due to the depolarized state of action potential. These effects were abolished by the addition of lidocaine (10 mg/L) to NL. We conclude that electrotonic interactions at NL-J depolarize the transmembrane potential and facilitate automaticity and triggered activity. Therefore, the NL-J site acted as a pacemaker zone capable of inducing arrhythmias which were abolished by lidocaine.

**Acidosis Influences ATP-Regulated  $\text{K}^+$  Channels in Normal and Hypertrophied Left Ventricular Myocytes**

Javier Cuevas, B.A., Shinichi Kimura, M.D., John S. Camero, Ph.D., Tetsushi Furukawa, M.D., Robert J. Myerburg, M.D., F.A.C.C., Arthur L. Bassett, Ph.D., University of Miami School of Medicine, Miami FL.

Cardiac  $\text{K}^+$ -ATP channels, activated by decreases in intracellular [ATP], may play a role in action potential shortening during ischemia. Patch-clamping was used to determine the effects of altering pH on properties of  $\text{K}^+$ -ATP channels in single endocardial myocytes isolated from normal and hypertrophied (via renal hypertension) cat LV.  $\text{K}^+$ -ATP channels of inside-out patches were bilaterally exposed to 140mM  $\text{K}^+$  solution (pH ranging from 7.4 to 6.0;  $22 \pm 1^\circ\text{C}$ ) in the absence of ATP. At pH 7.4,  $\text{K}^+$ -ATP channels from both normal and hypertrophied cells demonstrated similar slope conductances ( $\sim 65$  pS) during hyperpolarization, while channels from normal cells showed greater inward rectification during strong depolarizations. 80mV depolarizations at pH 7.4 produced significantly greater outward conductance ( $56 \pm 4$  pS,  $p < 0.05$ ) and higher open state probability ( $0.42 \pm 0.11$ ,  $p < 0.02$ ) in the hypertrophied myocyte  $\text{K}^+$ -ATP channel relative to normals ( $51 \pm 2$  pS and  $0.29 \pm 0.36$ ). Decreases in conductance and open-state probability occurred during acidosis in channels from both cell types. However, as external (intracellular) pH was lowered from 7.4 to 6.0 the absolute decreases in hypertrophy  $\text{K}^+$ -ATP open-state probability ( $0.33 \pm 0.10$ ) and conductance ( $21 \pm 3$  pS) at  $+80$  mV were 23% and 55% greater than in normals ( $0.25 \pm 0.05$  and  $14 \pm 1$  pS). These data show that ischemia-associated decreases in  $\text{pH}_i$  may modulate  $\text{K}^+$ -ATP channel properties and cellular action potential responses in normal and diseased myocardium.

**CELLULAR ELECTRICAL UNCOUPLING, CONTRACTURE AND CONDUCTION DURING THE EARLY PHASE OF MYOCARDIAL ISCHEMIA: DISCORDANT EFFECTS OF VERAPAMIL OR HYPOCALCEMIA.**

Wayne E. Cascio, M.D., Yan Gen-Xin, M.S., André G. Kléber, M.D., Department of Physiology, University of Bern, Switzerland

Both verapamil (V) and hypocalcemia (low  $[\text{Ca}^{2+}]_o$ ) diminish cardiac contractility, yet have dissimilar effects on action potential duration (APD) and the resting membrane potential (RMP). The purpose of this study was to contrast the effects of V and low  $[\text{Ca}^{2+}]_o$  on the magnitude and time course of changes in electrical and mechanical properties of myocardium during the early phase of ischemia. Arterially perfused isolated rabbit papillary muscles were suspended in a  $\text{H}_2\text{O}$ -saturated atmosphere. Standard intra- and extracellular electrodes, and a piezo-resistive element were used to measure simultaneously action potential characteristics, passive electrical properties (extracellular longitudinal resistance,  $r_o$ , and the intracellular longitudinal resistance,  $r_i$ ), conduction velocity ( $\Theta$ ), isometric twitch tension and resting tension. Three conditions were studied before and during no-flow ischemia: (1) perfusion with normal blood perfusate ( $n=12$ , extracellular ionized calcium concentration,  $[\text{Ca}^{2+}]_o = 1.0$  mM), (2) verapamil (0.5  $\mu\text{M}$ ,  $n=14$ ) normal blood perfusate, or (3) low  $[\text{Ca}^{2+}]_o$  blood perfusate ( $n=9$ ,  $[\text{Ca}^{2+}]_o = 0.4$  mM).

Perfusion for 30min with V or a low  $[\text{Ca}^{2+}]_o$  resulted in depolarization of the resting membrane potential (RMP,  $3 \pm 1$  mV and  $7 \pm 2$  mV, respectively), a decrease of the  $r_o$  ( $-15 \pm 4\%$  and  $-26 \pm 7\%$ ); and an increase in  $\Theta$  ( $4 \pm 2\%$  and  $6 \pm 2\%$ ). Neither intervention effected  $r_i$ . In contrast, V decreased ( $-20 \pm 6$  ms) and low  $[\text{Ca}^{2+}]_o$  increased ( $37 \pm 5$  ms) the APD at the time of 50% of repolarization (APD<sub>50</sub>). During the early phase of ischemia, V and low  $[\text{Ca}^{2+}]_o$  showed discordant effects despite large pre-ischemic reductions in the isometric twitch tension ( $48 \pm 4\%$  and  $78 \pm 3\%$ , respectively). After 8min of ischemia the following values and changes ( $\Delta$ ) relative to the pre-ischemic values were obtained: (\* indicates  $p < 0.05$  compared to control)

	$\Theta$ (cm/s)	$\Delta r_i$ (%)	$\Delta$ Resting Tension(mN)	RMP(mV)	$\Delta$ APD <sub>50</sub> (%)
Control	$40 \pm 3$	$17 \pm 7$	$-0.08 \pm 0.2$	$-66 \pm 2$	$68 \pm 6$
Verapamil	$50 \pm 3^*$	$18 \pm 13$	$0.02 \pm 0.06$	$-70 \pm 2$	$56 \pm 4^*$
Low $[\text{Ca}^{2+}]_o$	$40 \pm 4$	$87 \pm 32^*$	$0.20 \pm 0.09^*$	$-58 \pm 3^*$	$94 \pm 7^*$

These discordant effects of verapamil or low  $[\text{Ca}^{2+}]_o$  on cellular electrical coupling and ischemic contracture during the early phase of myocardial ischemia may in part be related to cellular calcium shifts influenced by APD and RMP.